

CORRESPONDENCE

Impact of Tumor Size on the Clinical Outcomes of Patients with Robson Stage I Renal Cell Carcinoma

I read the report by Kinouchi et al.,¹ in which they concluded that the tumor grading of renal cell carcinoma was not predictive of the survival of Robson Stage I patients. This does not mean that the three-grading system is worthless; the authors simply showed that the grading system of the Japanese General Rule for Clinical and Pathological Studies on Renal Cell Carcinoma² is not useful in the study of Robson Stage I renal cell carcinoma. The Japanese Rule² states that renal cell carcinoma should be classified according to the quantitatively dominant grade. Kinouchi et al.¹ omitted this explanation from their article. This "dominant rule" is obviously distinct from the other grading systems, according to which renal tumor grading is decided by the highest grade.^{3,4} I am afraid that their report might mislead international readers. The Japanese General Rule for Clinical and Pathological Studies on Renal Cell Carcinoma² was made in 1992. Its classification is different from "modern classification"⁵⁻⁷ and even from the classification criteria of the Armed Forces Institute of Pathology.³ Kinouchi et al.¹ nicely demonstrated the limitation of the Japanese Rule.² I would like to announce that it is time for the Japanese to throw away their old-fashioned Rule.² The recently published protocol of the Cancer Committee of the College of American Pathologists⁸ has greater possibilities.

REFERENCES

1. Kinouchi T, Saiki S, Meguro N, Maeda O, Kuroda M, Usami M, et al. Impact of tumor size on the clinical outcomes of patients with Robson Stage I renal cell carcinoma. *Cancer* 1999;85:689-95.
2. Japanese Urological Association, the Japanese Society of Pathology, Japan Radiological Society. General rule for clinical and pathological studies on renal cell carcinoma. 2nd edition. Tokyo: Kanehara Co. Ltd., 1992.
3. Murphy WM, Beckwith JB, Farrow GM. Tumors of the adult kidney. In: Rosai J, Sobin LH, eds. Tumors of the kidney, bladder, and related urinary structures. Atlas of tumor pathology. 3rd series, Fascicle 11. Washington, DC: Armed Forces Institute of Pathology, 1994:92-125.
4. Medeiros LJ, Jones EC, Alzawa S, Aldape HC, Chevillie JC, Goldstein NS, et al. Grading of renal cell carcinoma: Workgroup No. 2. Union Internationale Contre le Cancer (UICC) and the American Joint Committee on Cancer (AJCC). *Cancer* 1997;80:990-1.
5. Thoenes W, Störkel S, Rumpelt HJ. Histopathology and classification of renal cell tumors (adenomas, oncocytomas and carcinomas): the basic cytological and histopathological elements and their use for diagnostics. *Pathol Res Pract* 1986;181:125-43.
6. Kovacs G, Akhtar M, Beckwith BJ, Bugert P, Cooper CS, Delahunt B, et al. The Heidelberg classification of renal cell tumours [editorial]. *J Pathol* 1997;183: 131-3.
7. Störkel S, Eble JN, Adlakha K, Amin M, Blute ML, Bostwick DG, et al. Classification of renal cell carcinoma: Workgroup No. 1. Union Internationale Contre le Cancer (UICC) and the American Joint Committee on Cancer (AJCC). *Cancer* 1997;80:987-9.

8. Farrow G, Amlin MB. Protocol for the examination of specimens from patients with carcinomas of renal tubular origin, exclusive of Wilms tumor and tumors of urothelial origin: a basis for checklists. Cancer Committee, College of American Pathologists. *Arch Pathol Lab Med* 1999;123:23-7.

Naoki Kanomata, M.D.
Department of Pathology
Hyogo Prefectural Kakogawa Hospital
Kakogawa, Hyogo, Japan

Author Reply

First of all, it appears that Dr. Kanomata has misunderstood the grading system of the Japanese General Rule for Clinical and Pathological Studies on Renal Cell Carcinoma, which consists of a 3-stage system (i.e., Grades 1, 2, and 3). Apparently, Dr. Kanomata has misunderstood that in the Japanese General Rule, renal cell carcinoma should be classified according to the quantitatively dominant grade; however, the Japanese General Rule does not observe the grading system Dr. Kanomata described. For example, if tumors contain heterogenous grading patterns, the Japanese General Rule recommends describing the grading as G1>G2 when the major area contains G1 tumor and the minor area contains G2 tumor. If half of the tumor tissue samples contain G1 tumor and another half G2 tumor, the Japanese General Rule recommends describing the grading as G1=G2.

Second, we indicated in our article that higher grade determines the grading system, which is reasonable for predicting the patient's prognosis, as Dr. Kanomata mentioned.

Finally, we recommend that the Japanese General Rule determine the grading system according to the higher grade used.

Toshiaki Kinouchi, M.D., Ph.D.
Toshihiko Kotake, M.D.
Department of Urology
Osaka Medical Center for Cancer and
Cardiovascular Diseases
Osaka, Japan

Intraleural Administration of Interleukin-2 for the Treatment of Patients with Malignant Pleural Mesothelioma

A Phase II Study

We read with interest the article by Astoul et al.¹ regarding the administration of interleukin-2 (IL-2) for the treatment of patients with malignant mesothelioma. Although this appeared to be a well-designed and adequately conducted Phase II study, which included 22 patients who all received their planned therapy, we are concerned about the statistical analysis and interpretation of the results. Specifically, the authors compared the survival rate of responders to therapy with that of nonresponders, remarking that "... the median survival time of responders differed significantly from that of nonresponders: 28 months (SE: 12.12) and 8 months (SE: 5.07), respectively ($P < 0.01$)" and concluded that "these results confirm that IL-2 given intrapleurally has antitumor activity." A considerable amount has been written, however, about the pitfalls of analyses comparing responders to nonresponders (see, for example, Anderson et al.,² Weiss et al.,³ and Green et al.⁴). The main problem with this analysis is that patients have to live long enough to achieve a response; this "guarantee time" therefore biases the comparison in favor of the responders. Response status might also simply be a marker of patients with a better pretreatment prognosis who would have lived longer even had they not received therapy. Andersen et al.² discuss modifications of analyses based on response status that are less prone to bias, such as landmark analysis; however, none of these techniques can reliably be used to draw causal inferences regarding the effectiveness of treatment from Phase II data. Reviewers, authors, and editors must be aware of these important statistical issues as they generate medical literature.

REFERENCES

1. Astoul P, Picat-Joossen D, Viallat J-R, Boutin C. Intrapleural administration of interleukin-2 for the treatment of patients with malignant pleural mesothelioma: a Phase II study. *Cancer* 1998;83:2099-104.
2. Anderson JR, Cain KC, Gelber RD. Analysis of survival by tumor response. *J Clin Oncol* 1983;1:710-9.
3. Weiss GB, Bunce H III, Hokanson JA. Comparing survival by responders and nonresponders after treatment: a potential source of confusion in interpreting cancer clinical trials. *Control Clin Trials* 1983;4:43-52.

4. Green S, Benedetti J, Crowley J. Clinical trials in oncology. London: Chapman and Hall, 1997:155-8.

Theodore G. Karrison, Ph.D.
Nicholas J. Vogelzang, M.D.
*Departments of Health Studies and Medicine
University of Chicago
Chicago, Illinois*

Author Reply

We appreciate the comments of Dr. Karrison and Dr. Vogelzang regarding our article on intrapleural interleukin-2 (IL-2) therapy for patients with malignant pleural mesothelioma, and we thank them for their interest in reading our article.¹

When we statistically compared the survival rate of responders to therapy with that of nonresponders in a descriptive fashion, it was to point out our interest in performing "second-look" thoracoscopy on patients treated intrapleurally to assess their responses and we develop this point later. We are aware of the pitfalls of such analysis, although the survival of our patients was long enough to achieve a response based on an evaluation performed 36 days after the beginning of treatment (all eligible patients were assessable for response) and consequently all patients had "guarantee time."

We also agree that response status might simply be a marker of patients with a better pretreatment prognosis. However, except for three patients with Stage IA disease, the others had advanced cancer and represented homogenous cohort, and only one patient presented a fibrosarcomatous type that suggested a poor prognosis.² Taking into account the total protocol dose delivered, the toxicity experienced, and the patient population, we do not think that survival was associated with some basic underlying factor.

The landmark method compares a group of patients who have responded by a landmark time with another group that has failed to respond. It is quite different from a comparison of people who respond to therapy with those who do not respond to therapy.³

We did not suggest that longer survival for responders was due to intrapleural IL-2 therapy; this was not the aim of our Phase II study. Our last sentence stresses the need for Phase III comparative studies. However, based on computed tomography (CT) scan and "second-look" thoracoscopy, we think that the evaluation of antitumoral response was suitable. It is the first time, to our knowledge, that such an evaluation was performed after intrapleural treatment. Ya-

sumoto et al. reported antitumoral responses in patients with malignant pleural effusions due to lung carcinoma treated with intrapleural IL-2.⁴ However, those antitumoral responses were based on CT scan and pleural fluid as well as the disappearance of malignant cells, which is questionable.

In our Phase II study, it was noteworthy that the patients we considered to be responders (even thoracoscopy was not easy after intrapleural therapy) were those with prolonged survival. However, it is our opinion that the 43% 24-month and 31% 36-month survival rates of patients with Stage II malignant pleural mesothelioma (which is usually associated with poor prognosis) must be confirmed by comparative randomized Phase III studies to establish that the longer survival of responders was due to the treatment.

REFERENCES

1. Astoul P, Picat-Joossen D, Viallat JR, Boutin C. Intrapleural administration of interleukin-2 for the treatment of patients with malignant pleural mesothelioma: a Phase II study. *Cancer* 1998;83:2099-104.
2. Boutin C, Rey F, Gouvernet J, Viallat JR, Astout P, Ledoray V. Thoracoscopy in pleural malignant mesothelioma: a prospective study of 188 consecutive patients. *Cancer* 1993;73:394-404.
3. Anderson JR, Cain KC, Gelber RD, Gelman RS. Analysis and interpretation of the comparison of survival by treatment outcome variables in cancer clinical trials. *Cancer Treat Rep* 1985;69:1139-44.
4. Yasumoto K, Miyazaki K, Nagashima A, Ishida T, Kuda T, Yano T, et al. Induction of lymphokine-activated killer cells by intrapleural instillations of recombinant interleukin-2 in patients with malignant pleurisy due to lung cancer. *Cancer Res* 1987;47:2184-7.

Phillippe Astoul, M.D., Ph.D.
Christian Boutin, M.D.
*Department of Pulmonary Diseases
Hôpital de la Conception
University of the Mediterranean
Marseille, France*

Expression of p53 Protein and Resistance to Preoperative Chemotherapy in Locally Advanced Gastric Carcinoma

Cascinu et al. investigated the association between p53 status and response to preoperative chemotherapy in locally advanced gastric carcinoma.¹ Their study was based on determining the true extent of the tumor so that the p53 status could be analyzed cor-

rectly. In their study, tumor extent was evaluated with 2 completely different methods: laparotomy in 53% (16 of 30) of patients and computed tomography (CT) scan of the abdomen in 47% (14 of 30) of patients. The authors did not include any detailed information regarding the true disease stage of the patients prior to the administration of chemotherapy; they stated only that all patients had locally advanced disease.

As we all are aware, evaluation of response to chemotherapy is difficult in patients with locally advanced gastric carcinoma. The primary tumor site usually involves an ill-defined mass rather than an easily measurable lesion. Although upper gastrointestinal barium examination may determine the size of gastric lesions, it is ineffective in determining the depth of gastric wall involvement. Imaging modalities that can detect tumor invasion through the gastric wall into perigastric structures play a major role in staging.² CT, one of the most frequently used modalities, can detect only enlarged lymph nodes and predict direct invasion of adjacent structures by gastric carcinoma. However, due to its limited spatial resolution, CT cannot detect the depth of a tumor spread within the gastric wall and also is incapable of distinguishing fat plane obliteration caused by tumor invasion, inflammation, or cachexia. It is well known that normal size lymph nodes may contain tumor. This also explains the limited ability of CT to detect lymph node metastases.² In a study performed by McFee and Aust,³ CT was shown to have a sensitivity of 67% and a specificity of 61% for detecting lymphadenopathy in patients with gastric carcinoma and it also failed to show pancreatic invasion, with a sensitivity of 27%.

Endoscopic ultrasound (EUS) is a highly accurate method for the assessment of gastrointestinal malignancies. It has become the standard imaging modality for local staging of gastric carcinoma.⁴ The accuracy of EUS in determining the extent of primary tumor infiltration ranges from 67–97%.^{5–11} EUS is superior to CT in its ability to show the layers of the gastrointestinal wall, but is not as accurate as CT in the assessment of regional lymph nodes (N2) or distant metastases.² The combined results in 326 patients in the assessment of tumor extent showed that the accuracy of EUS was remarkably superior to CT (EUS: range, 82–92%, and CT: range, 11–43%). The accuracy of lymph node assessment by EUS and CT were found to range from 74–87% and from 25–51%, respectively.^{8,11,12} Conversely, laparotomy is a valuable method with which to detect occult intraabdominal metastatic disease. After subsequent laparotomy >25% of patients are reported to have been found to be understaged by CT scanning.¹³ However, laparotomy may not define the tumor extent in the gastric wall, especially in patients

with less advanced disease with N0 status, which may lead to difficulties in the evaluation of the stage.

CT and EUS are complementary rather than competitive. EUS is superior in locoregional staging and CT is preferred for the diagnosis of distant metastases. It is clear that EUS is an invaluable method in gastric staging and should be combined with CT scanning of the abdomen or with a more invasive method such as laparotomy in determining the true stage of disease. In the study by Cascinu et al.¹ the two different methods used for staging differ greatly in their sensitivity and specificity, which may result in the erroneous evaluation of disease stage as well as response rates. Consequently, this would affect response rates to preoperative chemotherapy and its association with p53 status.

REFERENCES

1. Cascinu S, Graziano P, Ferro BD, Staccioli MP, Ligi M, Carnevali A, et al. Expression of p53 protein and resistance to preoperative chemotherapy in locally advanced gastric carcinoma. *Cancer* 1996;83:1917–22.
2. Robert AH Jr., Judy Y, Vincent DM. Diagnosis and staging of gastric cancer. *Semin Oncol* 1996;23(3):325–35.
3. McFee AS, Aust IB. Gastric carcinoma and CAT scan. *Gastroenterology* 1981;80:196–8.
4. Tio TL, Schouwink MH, Clikot RJ, Tytgat GN. Preoperative TNM classification of gastric carcinoma by endosonography in comparison with the pathological TNM system: a prospective study of 72 cases. *Hepatogastroenterology* 1989;36:51–6.
5. Ohashi S, Nakazawa S, Yoshino J. Endoscopic ultrasonography in the assessment of invasive gastric cancer. *Scand J Gastroenterol* 1989;24:1039–48.
6. Akahoshi K, Misawa T, Fujishima H, Chijiwa Y, Maruoka A, Okkubo A, et al. Preoperative evaluation of gastric cancer by endoscopic ultrasound. *Gut* 1991;32:479–82.
7. Botet JP, Lightdale CJ, Zauber AG, Gerdes H, Winawer SJ, Urmacher C, et al. Preoperative staging of gastric cancer: comparison of endoscopic US and dynamic CT. *Radiology* 1991;181:426–32.
8. Caletti G, Ferrari A, Brocchi B, Barbara L. Accuracy of endoscopic ultrasonography in the diagnosis and staging of gastric cancer and lymphoma. *Surgery* 1993;113:14–27.
9. Grimm H, Blumhoeller KP, Hamper K, Koch J, Henne-Bruns D, Soehendra N. Endosonography for preoperative locoregional staging of esophageal and gastric cancer. *Endoscopy* 1993;25:224–30.
10. Ziegler K, Sanft C, Zimmer T, Zeltz M, Pelsenberg D, Stein H, et al. Comparison of computed tomography, endosonography, and intraoperative assessment in TN staging of gastric carcinoma. *Gut* 1993;34:604–10.
11. Grimm H, Hamper K, Maydeo A. Accuracy of endoscopic ultrasound and computed tomography in determining local/regional spread of gastric cancer: results of a prospective controlled study [abstract]. *Gastrointest Endosc* 1991;37:279.
12. Ajani JA, Mansfield PE, Ota DM. Potentially resectable gastric carcinoma: current approaches to staging and preoperative therapy. *World J Surg* 1995;19:216–20.

13. Rösch T. Endosonographic staging of esophageal cancer: a review of literature results. *Gastrointest Endosc Clin N Am* 1995;5:537-47.

Mustafa Özgüroglu, M.D.
Gökhan Demir, M.D.
Department of Internal Medicine
Medical Oncology Section
Cerrahpasa Medical School
Istanbul University
Istanbul, Turkey

Author Reply

In their correspondence, Drs. Özgüroglu and Demir point out the major role of endoscopic ultrasound (EUS) in the staging of locally advanced gastric carcinoma. Given the lack of staging with BUS in our study, they question the reliability of our results regarding the predictive value of p53 expression for resistance to chemotherapy.¹

In our opinion, the criticisms by Drs. Özgüroglu and Demir are only partially convincing. First, some problems still are encountered with EUS imaging.²⁻⁴ Second, this examination may not be necessary for assessment of response to chemotherapy in the presence of unresectable disease with macroscopic locoregional tumor involvement defined by endoscopy plus biopsies and computed tomography (CT) scan.⁵

In some studies, EUS has displayed a tendency to overestimate tumor extent because of the coexistence of peritumoral inflammation, especially in the presence of ulcerated tumor. In addition, artifacts due to erroneous scanning (tangential) may simulate increased thickness of layers. BUS may have poor accuracy in determining surface extension of the tumor.²⁻⁴ Overestimation of the lymph node status may be observed due to inflammatory hypertrophy of lymph nodes being a source of error for EUS. Similar to CT imaging, identification by EUS of metastases to lymph nodes with normal size and microinfiltration may be very difficult. In general, figures regarding EUS accuracy for the determination of lymph node status may be lower than reported by Özgüroglu and Demir, with a range of 50-87%.²⁻⁴ Finally, recent data suggest that underestimation of lymph node status by BUS may be related to differences in the histologic subtype of gastric carcinoma, with a lower accuracy for the detection of lymph node metastasis from undifferentiated compared with differentiated tumors.^{6,7}

The definition of locally advanced gastric carcinoma may comprise different presentations of the disease. Patients may have high risk but resectable gastric carcinoma

or they may present with unresectable disease with large volume local spread.⁵ In our opinion, staging with BUS particularly is indicated for the selection of high risk patients with small volume disease who are candidates for surgery and are eligible for studies of neoadjuvant chemotherapy.³⁻⁵ In addition, recent experiences have suggested that staging with EUS may be predictive of the risk of postoperative recurrence after radical surgery for gastric carcinoma.⁷

In our series, patients underwent endoscopy with biopsies and CT scans that demonstrated unresectable disease with bulky regional involvement or clear signs of infiltration of the pancreas, aorta, omentum, esophagus, and liver. After chemotherapy, partial response was defined as evidence on both endoscopy and CT scan of a >50% reduction in the tumor or the complete disappearance but positive endoscopic biopsy at the site of the tumor. Complete response was defined as a negative CT scan together with normal endoscopic view with negative biopsies.

In the presence of conspicuous locoregional involvement, the imaging we adopted for defining tumor burden may be adequate.⁴ In addition, the major criteria for evaluating response reduced the risk of erroneous evaluation of tumor shrinkage.⁴ We believe the relation we found between p53 overexpression and resistance to chemotherapy to be reliable.

REFERENCES

- Casclu S, Graziano F, Del Ferro E, Staccioli MP, Ligi M, Carnevali A, et al. Expression of p53 protein and resistance to preoperative chemotherapy in locally advanced gastric carcinoma. *Cancer* 1998;83(9):1917-22.
- Pollack BJ, Chak A, Sivak MV. Endoscopic ultrasonography. *Semin Oncol* 1998;23:336-46.
- Wang JY, Hsieh JS, Huang YS, Huang CJ, Hou MF, Huang TJ. Endoscopic ultrasonography for preoperative locoregional staging and assessment of resectability in gastric cancer. *Clin Imaging* 1998;22:355-9.
- Rösch T. Endoscopic techniques in gastrointestinal oncology. *Curr Opin Oncol* 1994;6:413-26.
- Rougier P, Lasser P, Ducreaux M, Mahjoubi M, Bognel C, Elias D. Preoperative chemotherapy of locally advanced gastric cancer. *Ann Oncol* 1994;5:59-68.
- Hamada S, Akahoshi K, Chijiwa Y, Nawata H, Sasaki I. Relationship between histological type and endosonographic detection of regional lymphnode metastases in gastric cancer. *Br J Radiol* 1997;70:697-702.
- Smith JW, Brennan MF, Botet JF, Gerdes H, Lightdale CJ. Preoperative endoscopic ultrasound can predict the risk of recurrence after operation for gastric carcinomas. *J Clin Oncol* 1993;11:2360-5.

Francesco Graziano, M.D.
Stefano Casclu, M.D.
Division of Medical Oncology
Hospital of Pesaro
Pesaro, Italy